

Thermal and kinematic corrections to the microwave background polarization induced by galaxy clusters along the line of sight

A. D. Challinor[★], M. T. Ford and A. N. Lasenby

Astrophysics Group, Cavendish Laboratory, Madingley Road, Cambridge, CB3 0HE, UK

5 February 2008

ABSTRACT

We derive analytic expressions for the leading-order corrections to the polarization induced in the cosmic microwave background (CMB) due to scattering off hot electrons in galaxy clusters along the line of sight. For a thermal distribution of electrons with kinetic temperature $k_{\text{B}}T_{\text{e}} \sim 10 \text{ keV}$ and bulk peculiar velocity $V \sim 1000 \text{ kms}^{-1}$, the dominant corrections to the polarization induced by the primordial CMB quadrupole and the cluster peculiar velocity arise from electron thermal motion and are at the level of ~ 10 per cent in each case, near the peak of the polarization signal. When more sensitive measurements become feasible, these effects will be significant for the determination of transverse peculiar velocities, and the value of the CMB quadrupole at the cluster redshift, via the cluster polarization route.

Key words: cosmic microwave background – cosmology: theory – galaxies: clusters: general – polarization.

1 INTRODUCTION

Compton scattering of cosmic microwave background photons off electrons in ionised intra-cluster gas leads to intensity distortions along the line of sight (Sunyaev & Zel’dovich 1972, 1980). The same scattering mechanism will also change the polarization state of the CMB if the radiation has a quadrupolar anisotropy in the average rest frame of the electrons. Such a quadrupole can arise from several physical effects, including: (i) transformation of a primordial quadrupole moment in the frame in which the CMB dipole vanishes (hereafter referred to as the CMB frame); (ii) aberration and Doppler effects in the transformation of the non-quadrupole moments of the primordial anisotropy due to electron bulk velocities; (iii) anisotropic scattering in the cluster; (iv) gravitational lensing from dynamic structures. The first three of these mechanisms were originally studied by Sunyaev & Zel’dovich (Sunyaev & Zel’dovich 1980; Zel’dovich & Sunyaev 1980), while the fourth was considered by Gibilisco (1997).

Although the small levels of polarization expected from cluster effects fall below current experimental sensitivities, the prospect of detection with future CMB experiments, such as the *Planck Surveyor* satellite, has fuelled a recent rise in interest in the topic. Audit & Simmons (1999) have (numerically) computed the frequency dependence of the

polarization arising from the transformation of the CMB monopole for a cold cluster, and Itoh, Nozawa & Kohyama (1999) derived an analytic expression for the frequency dependence of this polarization mechanism, including the first two corrections arising from thermal effects. The claims from these two groups that the original results of Sunyaev & Zel’dovich were in error were met with a paper from Sazonov & Sunyaev (1999), who confirmed the earlier Sunyaev–Zel’dovich analysis, and derived analytic results for the frequency dependence of the polarization arising from the CMB monopole and primordial quadrupole in the limit of a cold electron gas. Sazonov & Sunyaev (1999) also discussed the polarization effect due to prior anisotropic scattering (which is second-order in the optical depth, τ).

In this paper we reconsider the question of polarization generation from clusters, in the limit where second-order optical depth and gravitational lensing effects can be ignored. We use a convenient, covariant representation of the linear polarization, which simplifies the geometric interpretation of our results, to derive simple analytic expressions for the frequency dependence of the polarization signal. Our expressions include leading-order thermal and kinematic corrections to the results derived by Sazonov & Sunyaev (1999), and confirm their results in the limit of low electron temperatures and bulk velocities. Our results for the thermal corrections to the polarization arising from the electron bulk velocities differ significantly from those given by Itoh et al. (1999). We show that for typical cluster parameters, the leading-

[★] Email: A.D.Challinor@mrao.cam.ac.uk

order corrections to the polarization from the primordial quadrupole and the electron bulk velocity amount to reductions in these effects of ~ 10 per cent at the peak of their signals. These corrections should thus be considered in any future analysis of forthcoming CMB polarization data in the directions of clusters, when such detections are permitted by improvements in receiver sensitivity.

The paper is arranged as follows. In Section 2 we outline our covariant procedure for calculating the scattering term, which appears in the Boltzmann equation, for an unpolarized, but otherwise arbitrary, radiation field in the CMB frame. In Section 3, we present our results for the polarization signal arising from the individual multipoles of the primordial CMB intensity, and discuss their relative importance for a typical cluster. We end in Section 4 with our conclusions. We employ units with $c = h = 1$ throughout, and a $+$ $-$ $-$ signature for the spacetime metric.

2 POLARIZATION DUE TO ELECTRON SCATTERING OF AN ANISOTROPIC RADIATION FIELD

We consider observations of a diffuse radiation field by observers comoving with a timelike four-velocity field u^a . We adopt such a Lagrangian (or 1+3 covariant) approach since it provides arguably the most transparent description of the physics. For a recent review of the 1+3 covariant approach in the context of relativistic cosmology see Ellis (1998). Relative to u^a , a photon with four-momentum p^a is observed to have frequency ν and propagation direction e^a such that $p^a = \nu(u^a + e^a)$. The photon propagation direction e^a lies in the instantaneous rest space of the observer, so that $u^a e_a = 0$. We refer to geometric objects which are perpendicular to u^a on every index, such as e^a , as projected tensors. Since Thomson scattering does not generate circular polarization (Chandrasekhar 1950), we need only consider linearly polarized radiation here. The linear polarization state of radiation with frequency ν , which is propagating along e^a , is most conveniently described by the symmetric linear polarization tensor $\mathcal{P}_{ab}(\nu, e^c)$. The linear polarization is a trace-free, projected tensor which is also perpendicular to e^a . Following Thorne (1980), we refer to projected tensors which are also orthogonal to e^a as being transverse, and we denote the transverse, trace-free part of a rank-two tensor by the superscript TT , so that $\mathcal{P}_{ab} = [\mathcal{P}_{ab}]^{\text{TT}}$. It is useful to introduce the (screen) projection tensor $\mathcal{H}_{ab} \equiv g_{ab} - u_a u_b + e_a e_b$, where g_{ab} is the metric of spacetime. $\mathcal{H}_{ab}$ is a transverse tensor which projects perpendicular to u^a and e^a . Making use of $\mathcal{H}_{ab}$, the properties of $\mathcal{P}_{ab}(\nu, e^c)$ are summarised by

$$\mathcal{P}_{ab} = \mathcal{P}_{(ab)}, \quad \mathcal{H}^c_a \mathcal{P}_{bc} = \mathcal{P}_{ab}, \quad \mathcal{H}^{ab} \mathcal{P}_{ab} = 0, \quad (1)$$

where round brackets denote symmetrisation on the enclosed indices. The linear polarization tensor is related to the Stokes parameters $Q(\nu, e^c)$ and $U(\nu, e^c)$ via

$$\mathcal{P}_{11} = Q/2, \quad \mathcal{P}_{12} = \mathcal{P}_{21} = U/2, \quad \mathcal{P}_{22} = -Q/2, \quad (2)$$

where the components of $\mathcal{P}_{ab}$ are on a comoving orthonormal tetrad with the 3-direction aligned with e^a .

The generation of polarization through electron scattering is most simply described in the original rest-frame of the electron. Since we shall ultimately give our results in

the frame in which the CMB dipole vanishes, we will need the transformation laws for the radiation variables under changes of the velocity field u^a . If we consider an alternative four-velocity field $\tilde{u}^a$, this can be written in terms of observable quantities in the u^a frame as $\tilde{u}^a = \gamma(u^a + v^a)$. Here v^a is a projected (with respect to u^a) four-vector which describes the relative velocity of an observer comoving with $\tilde{u}^a$ with respect to the u^a frame. The quantity γ is the Lorentz factor associated with the relative motion, $\gamma = \tilde{u}^a u_a$ or equivalently $\gamma^{-2} = 1 + v^a v_a$. Similarly, we can express u^a in terms of quantities in the $\tilde{u}^a$ frame, so that $u^a = \gamma(\tilde{u}^a + \tilde{v}^a)$, where $\tilde{v}^a$ is the relative velocity in the $\tilde{u}^a$ frame. Introducing the projection tensor $h_{ab} = g_{ab} - u_a u_b$ in the u^a frame, and a similar projection in the $\tilde{u}^a$ frame, $\tilde{h}_{ab}$, we have $\tilde{v}^a = -\tilde{h}^a_b v^b / \gamma$. If we consider a given photon with four-momentum p^a , the frequency $\tilde{\nu}$ and propagation direction $\tilde{e}^a$ observed in the $\tilde{u}^a$ frame are expressed in terms of the quantities in the u^a frame via the Doppler and aberration formulae:

$$\tilde{\nu} = \gamma \nu (1 + e^a v_a), \quad (3)$$

$$\tilde{e}^a = [\gamma(1 + e^b v_b)]^{-1} (u^a + e^a) - \gamma(u^a + v^a). \quad (4)$$

For a given photon four-momentum p^a , the screen projection tensor in the $\tilde{u}^a$ frame, $\tilde{\mathcal{H}}_{ab}$, is related to $\mathcal{H}_{ab}$ by

$$\begin{aligned} \tilde{\mathcal{H}}_{ab} &= \mathcal{H}_{ab} - 2[\nu(1 + e^c v_c)]^{-1} p_{(a} \mathcal{H}_{b)d} v^d \\ &\quad + [\nu(1 + e^c v_c)]^{-2} p_a p_b \mathcal{H}_{d_1 d_2} v^{d_1} v^{d_2}, \end{aligned} \quad (5)$$

while the linear polarization tensor in the $\tilde{u}^a$ frame, $\tilde{\mathcal{P}}_{ab}(\tilde{\nu}, \tilde{e}^c)$, can be expressed in terms of the linear polarization in the u^a frame as

$$\tilde{\nu}^{-3} \tilde{\mathcal{P}}_{ab}(\tilde{\nu}, \tilde{e}^c) = \nu^{-3} \tilde{\mathcal{H}}_a^{d_1} \tilde{\mathcal{H}}_b^{d_2} \mathcal{P}_{d_1 d_2}(\nu, e^c). \quad (6)$$

This transformation law ensures that the degree of polarization $\mathcal{P}(\nu, e^c) = \sqrt{(2\mathcal{P}_{ab}\mathcal{P}^{ab})}/I(\nu, e^c)$, where $I(\nu, e^c)$ is the radiation intensity, is independent of frame for a given photon momentum.

The dynamics of the radiation is described by the collisional Boltzmann equation, which for the linear polarization takes the form

$$\mathcal{L}[\nu^{-3} \mathcal{P}_{ab}(\nu, e^c)] = K_{ab}(\nu, e^c), \quad (7)$$

where the Liouville operator acts along the photon path $x^a(\lambda)$, $p^a(\lambda)$ in phase space, with $p^a = dx^a/d\lambda$, according to

$$\mathcal{L}[\mathcal{P}_{ab}(\nu, e^c)] = \mathcal{H}_a^{d_1} \mathcal{H}_b^{d_2} p^e \nabla_e \mathcal{P}_{d_1 d_2}(\nu, e^c), \quad (8)$$

where ∇_a is the spacetime covariant derivative, and $K_{ab}(\nu, e^c)$ describes the scattering. For CMB photons scattering off free electrons in hot clusters, the average photon energy in the rest frame of the scattering electron is small compared to the electron rest mass, so that electron recoil can be neglected. Furthermore, the effects of induced scattering and Pauli blocking are negligible, so that the electron-radiation interaction can be approximated by classical Thomson scattering. In this limit, the scattering term in the rest frame of a beam of electrons (which has four-velocity $\tilde{u}^a$) with number density $\tilde{n}_e$, is given by (Challinor 1999)

$$\tilde{\nu}^2 \tilde{K}_{ab}(\tilde{\nu}, \tilde{e}^c) = \tilde{n}_e \sigma_T \left\{ -\tilde{\mathcal{P}}_{ab}(\tilde{\nu}, \tilde{e}^c) \right.$$

$$+ \frac{1}{10} [\tilde{I}_{ab}(\tilde{\nu})]^{\text{TT}} + \frac{3}{5} [\tilde{\mathcal{E}}_{ab}(\tilde{\nu})]^{\text{TT}} \}, \quad (9)$$

where σ_{T} is the Thomson cross section. Here, $I_{ab}(\nu)$ and $\mathcal{E}_{ab}(\nu)$ are projected (with respect to some general u^a) symmetric, trace-free (PSTF) tensors which describe the intensity and electric polarization quadrupoles respectively in a covariant manner (Challinor 1999). In equation (9), the tildes on $\tilde{I}_{ab}$ and $\tilde{\mathcal{E}}_{ab}$ denote that the intensity and polarization quadrupoles are evaluated in the rest-frame of the electron beam, and the transverse trace-free operations should be understood to refer to the $\tilde{u}^a$ frame also. For general ℓ , we define PSTF intensity multipoles

$$I_{A_\ell}(\nu) = \frac{1}{\Delta_\ell} \int d\Omega e_{\langle A_\ell \rangle} I(\nu, e^c), \quad (10)$$

where $\Delta_\ell \equiv 4\pi(-2)^\ell(\ell!)^2/(2\ell+1)!$ and $d\Omega$ is the element of solid angle of e^a , so that

$$I(\nu, e^c) = \sum_{\ell=0}^{\infty} I_{A_\ell}(\nu) e^{A_\ell}, \quad (11)$$

where A_ℓ is shorthand for the index string $a_1 \dots a_\ell$ and $e_{A_\ell} \equiv e_{a_1} \dots e_{a_\ell}$. Angle brackets denote the PSTF part on the enclosed indices. The electric polarization multipoles are defined by

$$\mathcal{E}_{A_\ell}(\nu) = \frac{1}{\Delta_\ell} \frac{2\ell(\ell-1)}{(\ell+1)(\ell+2)} \int d\Omega e_{\langle A_{\ell-2} \mathcal{P}_{a_{\ell-1}a_\ell} \rangle}(\nu, e^c), \quad (12)$$

for $\ell \geq 2$. A complete multipolar description of the linear polarization requires the introduction of magnetic multipoles also (Kamionkowski, Kosowsky & Stebbins 1997; Seljak & Zaldarriaga 1997; Challinor 1999), but we shall not need their definition here.

In this paper we shall only be concerned with the polarization induced by the cluster in the limit of low optical depth. Furthermore, we shall ignore the effects of any primordial polarization since these will be small compared to the effects of the intensity quadrupole. With these restrictions, only the intensity quadrupole prior to the interaction with the cluster need be retained on the right-hand side of equation (9). Transforming to the frame of an observer whose four-velocity is u^a , the scattering term becomes

$$K_{ab}(\nu, e^c) = \frac{1}{10} \tilde{n}_e \sigma_{\text{T}} \tilde{\nu}^{-2} \mathcal{H}_a^{d_1} \mathcal{H}_b^{d_2} [\tilde{I}_{d_1 d_2}(\tilde{\nu})]^{\text{TT}}, \quad (13)$$

where the transverse trace-free operation refers to the $\tilde{u}^a$ frame, and we have used the transformation law for the scattering term:

$$K_{ab}(\nu, e^c) = \mathcal{H}_a^{d_1} \mathcal{H}_b^{d_2} \tilde{K}_{d_1 d_2}(\tilde{\nu}, \tilde{e}^c). \quad (14)$$

We shall compute the scattering term in the CMB frame, which has four-velocity u^a , so that the intensity dipole $I_a(\nu)$ vanishes. First we calculate the intensity quadrupole in the rest frame of a beam of electrons, in terms of the multipoles $I_{A_\ell}(\nu)$ in the CMB frame. Making use of the invariance of $I(\nu, e^c)/\nu^3$, we have that

$$\begin{aligned} \tilde{I}_{ab}(\tilde{\nu}) &= \frac{15}{8\pi} \sum_{\ell=0}^{\infty} \int d\tilde{\Omega} \left\{ \frac{I_{C_\ell}[\gamma\tilde{\nu}(1+\tilde{e}^d\tilde{v}_d)]}{[\gamma(1+\tilde{e}^d\tilde{v}_d)]^{(\ell+3)}} \right. \\ &\quad \times (\tilde{e}^{c_1} - \tilde{v}^{c_1}) \dots (\tilde{e}^{c_\ell} - \tilde{v}^{c_\ell}) \tilde{e}_{\langle a} \tilde{e}_{b \rangle} \left. \right\}, \end{aligned} \quad (15)$$

where $\tilde{v}^a$ is the relative velocity of u^a with respect to $\tilde{u}^a$, so that $u^a = \gamma(\tilde{u}^a + \tilde{v}^a)$, and the integral is over solid angles of the projected directions $\tilde{e}^a$ in the $\tilde{u}^a$ frame. Next, we use this result for $\tilde{I}_{ab}(\tilde{\nu})$ in equation (13) to obtain the scattering term in the CMB frame due to a beam of electrons with number density $d\tilde{n}_e$ in its rest frame, and with velocity $v^a = -h_b^a \tilde{v}^b/\gamma$ relative to the CMB. Finally, we express $d\tilde{n}_e$ in terms of the number density of the beam in the CMB frame, using $dn_e = \gamma d\tilde{n}_e$, and integrate over the electron distribution. Assuming a non-degenerate, thermal distribution of electrons at temperature T_e in the frame in which the electron bulk velocity vanishes, we have, in the CMB frame,

$$dn_e = \frac{n_e d^3 p_e}{4\pi m_e^3 \Theta_e K_2(1/\Theta_e)} \exp[-\gamma\Gamma(1+V^a v_a)/\Theta_e], \quad (16)$$

where V^a is the projected, relative bulk velocity, Γ is the associated Lorentz factor, and $d^3 p_e$ is the element of 3-momentum. Here, $\Theta_e \equiv k_B T_e/m_e$ is the dimensionless electron temperature, m_e is the electron mass, $K_2(x)$ is a modified Bessel function, and n_e is the electron number density in the frame in which the bulk velocity vanishes. Note that the electron number density in the CMB frame is Γn_e .

3 POLARIZATION FROM THE MULTIPOLES $I_{A_\ell}(\nu)$

In this section we compute the contribution to the scattering term $K_{ab}(\nu, e^c)$ from each primordial multipole $I_{A_\ell}(\nu)$ in turn, expressing our results as expansions in the small parameters $\Theta_e = k_B T_e/(511 \text{ keV})$ and $V \equiv \sqrt{-V^a V_a}$.

In linear perturbation theory, where there is no non-linear scattering, there are no spectral distortions in the primordial radiation, so that

$$I(\nu, e^c) = \frac{2\nu^3}{e^{\nu/[k_B \bar{T}_r(1+\delta_T)]} - 1}, \quad (17)$$

where $\bar{T}_r$ is the all-sky average CMB temperature, and $\delta_T(e^c)$ is the dimensionless temperature anisotropy. If we expand $\delta_T(e^c)$ in PSTF multipoles τ_{A_ℓ} (Maartens, Ellis & Stoeger 1995),

$$\delta_T(e^c) = \sum_{\ell=2}^{\infty} \tau_{A_\ell} e^{A_\ell}, \quad (18)$$

(recall there is no dipole in the CMB frame), then, to first-order in the primordial anisotropy, we have for $\ell > 1$

$$I_{A_\ell}(\nu) = i(x) \tau_{A_\ell}, \quad \text{where,} \quad i(x) \equiv I(\nu) \frac{x e^x}{e^x - 1}, \quad (19)$$

with $I(\nu)$ the intensity monopole (a Planck spectrum with temperature $\bar{T}_r$), and $x \equiv \nu/(k_B \bar{T}_r)$ the dimensionless frequency. In this manner, the frequency dependence of the intensity multipoles is contained in the scalar $i(x)$ only, which proves to be very convenient for subsequent calculations. It should be noted that since $\delta_T(e^c)$ has no monopole component by definition, no confusion should arise from our later use of the symbol τ for the optical depth through the cluster.

3.1 The monopole contribution

The contribution to the intensity quadrupole in the rest frame of the electron beam from the intensity monopole in the CMB frame, $I(\nu)$, follows from equation (15):

$$\tilde{I}_{ab}(\tilde{\nu}) = \frac{15}{8\pi} \int d\tilde{\Omega} \frac{I[\gamma\tilde{\nu}(1 + \tilde{e}^c\tilde{v}_c)]}{[\gamma(1 + \tilde{e}^c\tilde{v}_c)]^3} \tilde{e}_{(a}\tilde{e}_{b)}, \quad (20)$$

which evaluates to

$$\tilde{I}_{ab}(\tilde{\nu}) = \frac{15}{8\beta^2} \int_{-1}^{+1} dx \frac{I[\gamma\tilde{\nu}(1 - \beta x)]}{[\gamma(1 - \beta x)]^3} (3x^2 - 1) \tilde{v}_{(a}\tilde{v}_{b)}, \quad (21)$$

where $\beta^2 = -\tilde{v}^a\tilde{v}_a = -v^a v_a$. Note that the contribution to $\tilde{I}_{ab}(\tilde{\nu})$ from the CMB monopole is $O(\beta^2)$. To transform to the CMB frame, we require the result

$$\mathcal{H}_a^{c_1} \mathcal{H}_b^{c_2} [\tilde{v}_{(c_1}\tilde{v}_{c_2)}]^{TT} = [\gamma(1 + e^c v_c)]^{-2} [v_{(a} v_{b)}]^{TT}, \quad (22)$$

which follows from elementary manipulations. The transverse trace-free and PSTF operations on the left-hand side of equation (22) refer to the $\tilde{u}^a$ frame, while those on the right refer to the u^a frame. It follows that the scattering term $K_{ab}(\nu, e^c)$ due to a beam of electrons with relative velocity v^a is proportional to $[v_{(a} v_{b)}]^{TT}$, so that the radiation is polarized along the normal to the plane containing v^a and the line of sight e^a .

We use equations (21) and (22) in the right-hand side of equation (13), and expand in powers of β . The integral over the electron distribution is handled by an expansion in powers of V and Θ_e . The integrals are straightforward, but tedious, making them ideally suited to symbolic computer algebra packages (we use MAPLE). The final result is an asymptotic expansion for the scattering term in the CMB frame due to the intensity monopole, of which the leading few terms are

$$\begin{aligned} \nu^2 K_{ab}(\nu, e^c) = & \frac{1}{10} n_e \sigma_T [V_{(a} V_{b)}]^{TT} I(\nu) \frac{x e^x}{e^x - 1} \left\{ \frac{1}{2} F \right. \\ & + \Theta_e \left[3F - 2(2F^2 + G^2) \right. \\ & + \left. \left. \frac{1}{2} F(F^2 + 2G^2) \right] \right. \\ & + \left. \left. V\mu \left[-\frac{1}{2} F + \frac{1}{4} (2F^2 + G^2) \right] \right\}, \quad (23) \end{aligned}$$

where $\mu \equiv -V^a e_a / V$ is the cosine of the angle between the bulk velocity of the cluster and the photon propagation direction. Here, $F \equiv x \coth(x/2)$ and $G \equiv x / \sinh(x/2)$. The presence of the factor $[V_{(a} V_{b)}]^{TT}$ ensures that the radiation is polarized along the normal to the plane containing the bulk velocity and the line of sight, as required by symmetry considerations.

If we assume that the distance across the cluster is small compared to the characteristic length scale of the local spacetime curvature, then in the limit of low optical depth the emergent polarization is simply $\mathcal{P}_{ab}(\nu, e^c) = \nu^2 K_{ab}(\nu, e^c) t$, where t is the light travel time across the cluster in the CMB frame. To obtain the degree of polarization $\mathcal{P}(\nu, e^c)$ from equation (23), one should multiply by t , take the magnitude (times $\sqrt{2}$), and divide by the emergent intensity $I(\nu, e^c) + \Delta I(\nu, e^c)$, where $\Delta I(\nu, e^c)$ is the change in intensity due to the scattering. Ignoring the small anisotropy in the CMB frame, the leading order terms of $\Delta I(\nu, e^c)$ are (Sunyaev & Zel'dovich 1972; Sunyaev & Zel'dovich 1980)

$$\Delta I(\nu, e^c) = n_e \sigma_T t I(\nu) \frac{x e^x}{e^x - 1} [\Theta_e(F - 4) + V\mu]. \quad (24)$$

Higher order terms in the expansion of $\Delta I(\nu, e^c)$ have been calculated recently (Stebbins 1997; Challinor & Lasenby 1998; Challinor & Lasenby 1999; Itoh, Kohyama & Nozawa 1998; Nozawa, Itoh & Kohyama 1998; Sazonov & Sunyaev 1998). Identifying the effective optical depth through the cluster, $\tau \equiv n_e \sigma_T t$, in the limit of small Θ_e and V we recover the result for $\mathcal{P}(\nu, e^c)$ given as equation (22) in Sazonov & Sunyaev (1999):

$$\mathcal{P}(\nu, e^c) = \frac{x^2 e^x}{20(e^x - 1)} \coth(x/2) \tau V_t^2, \quad (25)$$

where we have used

$$2[V_{(a} V_{b)}]^{TT} [V^{(a} V^{b)}]^{TT} = V_t^4, \quad (26)$$

with $V_t^2 = V^2(1 - \mu^2)$ the square of the transverse velocity.

The $O(\Theta_e^0, V^0)$ and $O(\Theta_e^1, V^0)$ terms in the curly braces on the right-hand side of equation (23) correct those given by Itoh et al. (1999). Itoh et al. also give an expression for the $O(\Theta_e^2, V^0)$ term; we include our result here for completeness:

$$\begin{aligned} \Theta_e^2 \left\{ \frac{15}{2} F - 21(2F^2 + G^2) + \frac{867}{28} F(F^2 + 2G^2) \right. \\ - \frac{40}{7} \left[F^4 + \frac{1}{2} G^2(11F^2 + 2G^2) \right] \\ + \left. \frac{2}{7} F \left[F^4 + \frac{1}{2} G^2(26F^2 + 17G^2) \right] \right\}. \quad (27) \end{aligned}$$

In the Rayleigh-Jeans region, neglecting $\Delta I(\nu, e^c)$ compared to $I(\nu, e^c) \approx I(\nu)$, we find the degree of polarization

$$\mathcal{P}(\nu, e^c) = \frac{1}{10} \tau V_t^2 \left(1 - 6\Theta_e + 2V\mu + \frac{183}{7} \Theta_e^2 + \dots \right), \quad (28)$$

which, as noted by Sazonov & Sunyaev (1999), recovers the result of Sunyaev & Zel'dovich (1980) at lowest order. The $O(\Theta_e^0, V^1)$ term in the braces on the right of equation (23) is responsible for the weak dependence of the degree of polarization on the absolute value of the bulk velocity, for a given transverse velocity V_t , as found by Audit & Simmons (1999) through numerical calculations in the zero temperature limit.

In Fig. 1 we plot the frequency dependence of the magnitude of the polarization, $I(\nu)\mathcal{P}(\nu, e^c)$, in units of $2(k_B T_c)^3 / (hc)^2$, for the cluster parameters $k_B T_e = 10$ keV, $V_t = 1000 \text{ km s}^{-1}$, $\tau = 0.01$. The direction of the bulk motion is chosen to maximise the $O(\Theta_e^0, V^1)$ term in equation (23), so that $\mu = 1/\sqrt{3}$. For these parameters, the leading order correction to the polarization from the CMB monopole is thermal in origin, and tends to broaden the polarization signal. In the Rayleigh-Jeans region the correction amounts to a reduction in the signal of $\simeq 10$ per cent, which also follows from equation (28), while near the peak of the polarized signal ($x = 4.85$) the reduction is $\simeq 15$ per cent. Near the peak, including the $O(\Theta_e^2, V^0)$ term in curly braces in equation (23) gives a further 3 per cent correction upwards, while including the $O(\Theta_e^0, V^0)$ term (the leading order kinematic correction) gives a further correction upwards of only 1 per cent when the radial motion is towards the cluster. In the Wien region, where the degree of polarization is large (Audit & Simmons 1999), thermal effects lead to a much larger fractional enhancement in the polarization. For example, in the *Planck* HFI channel at 545 GHz, the thermal enhancement of the polarization due to the monopole is nearly 30




Figure 1. Magnitude of the polarization [in units of $2(k_B \bar{T}_e)^3/(hc)^2$] due to the CMB monopole for a cluster with $k_B T_e = 10$ keV, $V_t = 1000$ km s $^{-1}$, $\tau = 0.01$, and $\mu = 1/\sqrt{3}$. The dashed-dotted line is the contribution with only the $O(\Theta_e^0, V^0)$ term in curly braces in equation (23) included. The solid line also includes the $O(\Theta_e^1, V^0)$ term, and the dashed line further includes the $O(\Theta_e^2, V^0)$ term given as equation (27). The dotted line includes these first two thermal corrections, as well as the $O(\Theta_e^0, V^1)$ term.

per cent. However, for all of the *Planck* channels, the degree of polarization from the cluster parameters in Fig. 1 falls well below the predicted detector sensitivity. In the *Planck* HFI channels at 143, 217 and 545 GHz, the magnitude of the polarization from the monopole as an equivalent fractional temperature change $\delta T/T$ is $\simeq 0.014$, 0.019 , and 0.070 μ K/K respectively. These values should be compared to the predicted sensitivities of 3.7, 8.9 and 208 μ K/K. Our conclusions concerning the feasibility of direct detection of cluster polarization with the *Planck* satellite are less optimistic than those in Audit & Simmons (1999). This is due to our use of rather more realistic cluster parameters, and the correct predicted equivalent thermodynamic temperature sensitivities.

The errors in our analytic expressions for the polarization (and also the intensity change) decrease with decreasing cluster temperature and bulk velocity. For fixed parameters, the errors increase with increasing frequency. The latter behaviour is due to the use of a Fokker-Planck expansion of the intensity with respect to the frequency change on scattering in the CMB frame, during the evaluation of $\tilde{I}_{ab}(\tilde{\nu})$ [equation (15)]. In practice this limits the range of frequencies for which the expansions are valid to $x \lesssim 1/\beta$ (recall that β is the speed of the scattering electron relative to the CMB). For a typical cluster, the thermal electron motion is the limiting factor, giving a maximum dimensionless frequency x of the order of 10. Note that equation (25) suggests a degree of polarization which behaves as x^2 for large x . However, this is not in conflict with the requirement that $\mathcal{P}(\nu, e^c)$ be bounded by unity given the limited range in x for which equation (25) is valid.

3.2 The quadrupole contribution

The contribution to the quadrupole in the rest frame of a beam of electrons from the primordial quadrupole in the

CMB frame is given by

$$\tilde{I}_{ab}(\tilde{\nu}) = \frac{15}{8\pi} \int d\tilde{\Omega} \left\{ \frac{i[\gamma\tilde{\nu}(1 + \tilde{e}^d \tilde{v}_d)]}{[\gamma(1 + \tilde{e}^d \tilde{v}_d)]^5} \times \tau_{c_1 c_2} (\tilde{e}^{c_1} - \tilde{v}^{c_1})(\tilde{e}^{c_2} - \tilde{v}^{c_2}) \tilde{e}_{\langle a} \tilde{e}_{b \rangle} \right\}. \quad (29)$$

Evaluating this integral is a little more involved than for the monopole calculation given in the previous subsection. An expansion in powers of β gives three distinct sets of terms. First, there are terms proportional to $\tilde{h}_{\langle a}^{c_1} \tilde{h}_{b \rangle}^{c_2} \tau_{c_1 c_2}$, where it will be recalled that $\tilde{h}_{ab} \equiv g_{ab} - \tilde{u}_a \tilde{u}_b$ is the projection tensor into the instantaneous rest space of the $\tilde{u}^a$ frame. These terms transform to the CMB frame according to

$$\mathcal{H}_a^{c_1} \mathcal{H}_b^{c_2} [\tilde{h}_{\langle c_1}^{d_1} \tilde{h}_{c_2 \rangle}^{d_2} \tau_{d_1 d_2}]^{\text{TT}} = [\tau_{ab}]^{\text{TT}} - \frac{2}{\alpha} [v_{\langle a} \tau_{b \rangle c} e^c]^{\text{TT}} + \frac{1}{\alpha^2} \tau_{c_1 c_2} e^{c_1} e^{c_2} [v_{\langle a} v_{b \rangle}]^{\text{TT}}, \quad (30)$$

where $\alpha \equiv 1 + e^c v_c$. Second, there are terms proportional to $\tilde{v}_{\langle a} \tilde{h}_{b \rangle}^{c_1} \tilde{v}^{c_2} \tau_{c_1 c_2}$, which transform as

$$\mathcal{H}_a^{c_1} \mathcal{H}_b^{c_2} [\tilde{v}_{\langle c_1}^{d_1} \tilde{h}_{c_2 \rangle}^{d_2} \tilde{v}^{d_2} \tau_{d_1 d_2}]^{\text{TT}} = -\frac{1}{\gamma \alpha} [v_{\langle a} \tau_{b \rangle c} v^c]^{\text{TT}} + \frac{1}{\gamma \alpha^2} [v_{\langle a} v_{b \rangle}]^{\text{TT}} \tau_{c_1 c_2} v^{c_1} e^{c_2}. \quad (31)$$

In equations (30) and (31) the transverse trace-free and PSTF operations on the left-hand sides refer to the $\tilde{u}^a$ frame, while those on the right refer to the u^a frame. Finally, there are terms proportional to $\tilde{v}_{\langle a} \tilde{v}_{b \rangle} v^{c_1} v^{c_2} \tau_{c_1 c_2}$, whose transformation law follows from equation (22). It follows that, in general, the polarization due to the primordial quadrupole will be a superposition of six geometric objects: the five terms on the right-hand sides of equations (30) and (31), with v^a replaced by the cluster peculiar velocity V^a , and a term $V_{\langle a} V_{b \rangle} V^{c_1} V^{c_2} \tau_{c_1 c_2}$. If the peculiar velocity of the cluster vanishes, the polarization tensor is proportional to $[\tau_{ab}]^{\text{TT}}$ only. The latter case was discussed by Sazonov & Sunyaev (1999) in the low temperature limit; it is straightforward to show that the dependence of the degree of polarization on the position of the cluster on the sky, for $\mathcal{P}_{ab}(\nu, e^c) \propto [\tau_{ab}]^{\text{TT}}$, is equivalent to that derived in Sazonov & Sunyaev (1999) by non-geometric means.

The leading few terms for the scattering term in the CMB frame due to the primordial quadrupole are

$$\nu^2 K_{ab}(\nu, e^c) = \frac{1}{10} n_e \sigma_T I(\nu) \frac{x e^x}{e^x - 1} \left([\tau_{ab}]^{\text{TT}} \left\{ 1 + \Theta_e [-6F + \frac{1}{2}(2F^2 + G^2)] + V\mu(-2 + F) \right\} - 2[V_{\langle a} \tau_{b \rangle c} e^c]^{\text{TT}} \right). \quad (32)$$

For completeness, we also give the $O(\Theta_e^2)$ term. This only has a $[\tau_{ab}]^{\text{TT}}$ geometric dependence, and so would appear in the curly braces in equation (32):

$$\Theta_e^2 \left\{ -15F + \frac{69}{4}(2F^2 + G^2) - \frac{64}{7}F(F^2 + 2G^2) + \frac{4}{7} \left[F^4 + \frac{1}{2}G^2(11F^2 + 2G^2) \right] \right\}. \quad (33)$$

In the limit of small Θ_e and V , the polarization is given by

$$\mathcal{P}_{ab}(\nu, e^c) = \frac{1}{10} \tau i(x) [\tau_{ab}]^{\text{TT}} = \frac{1}{10} \tau [I_{ab}(\nu)]^{\text{TT}}, \quad (34)$$




Figure 2. Magnitude of the dominant $[\tau_{ab}]^{\text{TT}}$ polarization [in units of $2(k_B \bar{T}_r)^3/(hc)^2$] due to the CMB quadrupole for a cluster with $k_B T_e = 10 \text{ keV}$, $V\mu = 1000 \text{ kms}^{-1}$ and $\tau = 0.01$. We use the best-fitting *COBE* results for the quadrupole: $\lambda_1 = -25 \mu\text{K}$, $\lambda_2 = 23 \mu\text{K}$ and $\lambda_3 = 2 \mu\text{K}$. The dashed-dotted line is the contribution with only the $O(\Theta_e^0, V^0)$ term in curly braces in equation (32) included. The solid line also includes the $O(\Theta_e^1, V^0)$ term, and the dashed line further includes the $O(\Theta_e^2, V^0)$ term given as equation (33). The dotted line includes these first two thermal corrections, as well as the $O(\Theta_e^0, V^1)$ term.

which also follows directly from equation (13) on setting $v^a = 0$. This very simple result is equivalent to that derived by Sazonov & Sunyaev (1999), but the presentation here benefits from the fully geometric approach. We can write τ_{ab} in the form $\bar{T}_r \tau_{ab} = \sum_{i=1}^3 \lambda_i n_a^{(i)} n_b^{(i)}$, where $\{\lambda_i\}$ are the (observed) eigenvalues of the temperature quadrupole, and $\{n_a^{(i)}\}$ are the corresponding unit projected eigenvectors. (For the metric signature employed here, $\bar{T}_r \tau_a^b n_b^{(i)} = -\lambda_i n_a^{(i)}$.) For $\mathcal{P}_{ab}(\nu, e^c) \propto [\tau_{ab}]^{\text{TT}}$, it follows that the degree of polarization is maximal along the eigendirection with intermediate eigenvalue (say λ_3). Perpendicular to this direction, the polarization vanishes for directions such that $(e^a n_a^{(1)})^2 = (\lambda_1 - \lambda_3)/(\lambda_1 - \lambda_2)$. For clusters at these positions on the sky, the polarization signal due to the primordial quadrupole will vanish except for small corrections arising from the cluster peculiar velocity. The first such correction is given in equation (32). Including only this correction, the degree of polarization due to the primordial quadrupole along directions for which $[\tau_{ab}]^{\text{TT}}$ vanishes evaluates to

$$\mathcal{P} = \frac{1}{10} \tau V_e 2[(\lambda_3 - \lambda_1)(\lambda_2 - \lambda_3)]^{1/2}, \quad (35)$$

which we have expressed as an equivalent temperature. In the Rayleigh-Jeans region, we find

$$\begin{aligned} \mathcal{P}_{ab}(\nu, e^c) &= \frac{1}{10} \tau I(\nu) [\tau_{ab}]^{\text{TT}} \left(1 - 6\Theta_e + \frac{183}{7} \Theta_e^2 \right) \\ &\quad - 2[V_{(a} \tau_{b)c} e^c]^{\text{TT}}. \end{aligned} \quad (36)$$

Note that the $O(V)$ correction to the $[\tau_{ab}]^{\text{TT}}$ polarization term vanishes in this limit.

In Fig. 2 we plot the magnitude of the dominant polarization signal (that with $[\tau_{ab}]^{\text{TT}}$ geometric dependence) due to the CMB quadrupole, for the cluster parameters $k_B T_e = 10 \text{ keV}$, $V\mu = 1000 \text{ kms}^{-1}$ and $\tau = 0.01$. For the CMB quadrupole we take the *COBE* best-fitting val-

ues $\lambda_1 = -25 \mu\text{K}$, $\lambda_2 = 23 \mu\text{K}$ and $\lambda_3 = 2 \mu\text{K}$ used by Sazonov & Sunyaev (1999). The line of sight is chosen to lie along $n_a^{(3)}$ to maximise the polarization signal. As with the polarization due to the CMB monopole, we see that the dominant correction to the polarization sourced by the primordial quadrupole is due to thermal effects, and these tend to broaden the signal. In the Rayleigh Jeans region the size of the fractional correction is the same as for the signal due to the monopole $\simeq 10$ per cent [compare equations (28) and (36)]. Near the signal peak ($x = 3.83$) the leading thermal correction leads to a reduction of $\simeq 14$ per cent, but this is reduced to $\simeq 12$ per cent when the next order thermal correction and the $O(V)$ kinematic term are included. The corrections are most significant in the Wien region; in the 545 GHz *Planck* HFI channel the thermal terms increase the signal by $\simeq 60$ percent. However, since the monopole contribution to the degree of polarization rises more rapidly with frequency (as x^2) than the quadrupole (as x) for large x , the quadrupole contribution is likely to be less significant than the monopole in the Wien region, for typical cluster parameters (Sazonov & Sunyaev 1999). In the *Planck* HFI channels at 143, 217 and 545 GHz, the magnitude of the polarization from the quadrupole as an equivalent fractional temperature change $\delta T/T$ is $\simeq 0.016$, 0.016 , and $0.029 \mu\text{K/K}$ respectively, which, like the monopole contribution, is well below the predicted sensitivities.

3.3 The contribution of higher multipoles

The ℓ -th multipole of the CMB anisotropy gives rise to a quadrupole of $O(\beta^{\ell-2})$ in the rest frame of the scattering electron. It follows that the leading terms in the signal due to the CMB octupole ($\ell = 3$) will be $O(V, \Theta_e)$. Unlike the monopole, the octupole can give a thermal polarization signal even if the cluster peculiar velocity vanishes ($V = 0$), due to the preferred directions in space that it defines. Since the octupole is of similar magnitude to the quadrupole [the *COBE* best-fitting values give $|\tau_{abc}|/|\tau_{ab}| \simeq 2.3$ (Stoeger, Araujo & Gebbie 1997)], and given that the $O(\Theta_e)$ term in the quadrupole contribution to the polarization can be a sizeable fraction of the total quadrupole signal, the octupole contribution should potentially be included in future attempts to determine the primordial quadrupole at the location of distant (hot) clusters via the cluster polarization route (Kamionkowski & Loeb 1997).

The calculation of the polarization due to the CMB octupole follows that of the monopole and quadrupole. For the leading two terms we find

$$\begin{aligned} \nu^2 K_{ab}(\nu, e^c) &= \frac{1}{10} n_e \sigma_T I(\nu) \frac{x e^x}{e^x - 1} \frac{3}{7} \left\{ [\tau_{abc} V^c]^{\text{TT}} (5 - F) \right. \\ &\quad \left. + \Theta_e [\tau_{abc} e^c]^{\text{TT}} \left[6F - \frac{1}{2} (2F^2 + G^2) \right] \right\}. \end{aligned} \quad (37)$$

For the hexadecapole ($\ell = 4$) the quadrupole in the rest frame of the electron is second-order in β , so we might anticipate an $O(\Theta_e)$ polarization signal. However, to this order the polarization from a beam of electrons with relative velocity v^a is proportional to $[\tau_{abcd} v^c v^d]^{\text{TT}}$, which results in a polarization signal proportional to $[\tau_{abcd} V^c V^d]^{\text{TT}}$ on integrating over the electron distribution. For reasonable values of the cluster peculiar velocity, we infer that the effects of

the CMB multipoles with $\ell > 3$ can be safely ignored, even for rather hot clusters.

4 CONCLUSION

We have presented a direct geometric method for determining the polarization due to the scattering of anisotropic radiation off a thermal distribution of electrons, with non-vanishing bulk velocity, in the limit of low optical depth. Our results for the polarization due to the CMB monopole and quadrupole fully agree with those obtained by Sazonov & Sunyaev (1999) [see also Zel'dovich & Sunyaev (1980)] in the low temperature limit. Our results for the thermal corrections to the monopole effect correct those given by Itoh et al. (1999). A careful inspection of their analysis suggests that the discrepancy arises from the omission of a rotation needed to relate the scattering basis in the rest frame of the initial electron and the CMB frame. We have shown that for typical cluster parameters the thermal corrections tend to dominate the kinematic corrections. Near the peak of their respective polarization signals, thermal effects tend to reduce the magnitude of the polarization from the monopole and quadrupole by $\simeq 12$ per cent. The corrections are much larger in the Wien part of the spectrum. We have also noted that the CMB octupole can lead to a polarization signal from hot clusters which is of similar magnitude to the thermal correction to the quadrupole signal. The effects reported here may be significant for future attempts to determine the CMB quadrupole at distant clusters and the tangential peculiar velocity, to an accuracy of a few per cent, via polarization observations.

ACKNOWLEDGEMENTS

ADC acknowledges Queens' College, Cambridge for support in the form of a Research Fellowship.

REFERENCES

- Audit E., Simmons J. F. L., 1999, preprint; astro-ph/9812310
 Challinor A. D., 1999, preprint; astro-ph/9903283
 Challinor A. D., Lasenby A. N., 1998, ApJ, 499, 1
 Challinor A. D., Lasenby A. N., 1999, ApJ, 510, 930
 Chandrasekhar S., 1950, Radiative Transfer. Clarendon Press, Oxford
 Ellis G. F. R., 1998, preprint; astro-ph/9812046
 Gibilisco M., 1997, Ap&SS, 249, 189
 Itoh N., Kohyama Y., Nozawa S., 1998, ApJ, 502, 7
 Itoh N., Nozawa S., Kohyama Y., 1999, preprint; astro-ph/9812376
 Kamionkowski M., Kosowsky A., Stebbins A., 1997, Phys. Rev. Lett., 78, 2058
 Kamionkowski M., Loeb A., 1997, Phys. Rev. D, 56, 4511
 Maartens R., Ellis G. F. R., Stoeger W. R., 1995, Phys. Rev. D, 51, 1525
 Nozawa S., Itoh N., Kohyama Y., 1998, ApJ, 508, 17
 Sazonov S. Y., Sunyaev R. A., 1998, ApJ, 508, 1
 Sazonov S. Y., Sunyaev R. A., 1999, preprint; astro-ph/9903287
 Seljak U., Zaldarriaga M., 1997, Phys. Rev. Lett., 78, 2054
 Stebbins A., 1997, preprint; astro-ph/9709065
 Stoeger W. R., Araujo M., Gebbie T., 1997, ApJ, 476, 435; see also astro-ph/9904346

- Sunyaev R. A., Zel'dovich Y. B., 1972, Comments Astrophys. Space Phys., 4, 173
 Sunyaev R. A., Zel'dovich Y. B., 1980, MNRAS, 190, 413
 Thorne K. S., 1980, Rev. Mod. Phys., 52, 299
 Zel'dovich Y. B., Sunyaev R. A., 1980, Astron. Lett., 6, 285

This paper has been produced using the Royal Astronomical Society/Blackwell Science L^AT_EX style file.